arXiv:1008.2572v2 [quant-ph] 11 Apr 2011

Entanglement in the ground state of interacting qubits driven by a quantum field

B. M. Rodríguez-Lara[1,2] and Ray-Kuang Lee[3,2]

[1]*Centre for Quantum Technologies, National University of Singapore,*
2 Science Drive 3, Singapore 117542.
[2]*Institute of Photonics Technologies,*
National Tsing-Hua University, Hsinchu 300, Taiwan.
[3]*Department of Physics, National Tsing-Hua University, Hsinchu 300, Taiwan.*

Abstract

We study the ground state entanglement properties of a finite size ensemble of interacting qubits driven by a quantized field. In the absence of driving, the system reduces to a Lipkin-Meshkov-Glick model that presents maximal entanglement at the first of a series of quantum phase transitions; *i.e.* a W-state of the qubit ensemble. We present exact and numerical solutions, within and without the rotating wave approximation, in that order, showing that a large finite probability of having the maximal entangled state in the ground state is preserved in the driven system.

Entanglement is a fundamental quantum mechanical phenomenon [1] and a precious resource in quantum information processing [2]. A multi-qubit W-state [3] maximizes pairwise entanglement of formation [4, 5] and is a robust source of entanglement [6, 7]; i.e., it retains maximal bipartite quantum correlations whenever any pair of qubits are traced out. Interaction between qubits is necessary to generate entangled states. For an ensemble of interacting qubits, the Lipkin-Meshkov-Glick (LMG) model is an example of a system producing maximal pairwise entanglement of formation at the phase transition of its ground state; *e.g.*, a collection of spins with (anti-)ferromagnetic interactions embedded in a magnetic field produce maximal entanglement at the (first)second order quantum phase transition [8].

Recently, it has been proposed that the LMG model, originally from nuclear physics, may describe the Josephson effect or two-mode Bose-Einstein condensate (BEC) [9]. Also, schemes to realize a dissipative LMG model in optical cavity-QED [10] and in circuit-QED [11] have been discussed.

In this work, we consider a system consisting of an ensemble of interacting qubits driven by a quantized field and described by the extended Dicke Hamiltonian,

$$\hat{H} = \omega_f \hat{N} + \delta \hat{J}_z + \eta N_a^{-1} \hat{J}_z^2 + \lambda N_a^{-1/2} \left(\hat{a} + \hat{a}^\dagger \right) \hat{J}_x, \tag{1}$$

which is equivalent to the LMG model in the limit $\lambda = 0$. The detuning δ has been defined as the difference between the qubit transition frequency ω and the quantized field frequency ω_f, $\delta = \omega - \omega_f$; the orbital angular momentum representation, J_i with $i = x, y, z, \pm$, has been chosen to describe the qubit ensemble, while the field is described by its creation (annihilation) operators, $\hat{a}$ ($\hat{a}^\dagger$) and the total excitation number is given by $\hat{N} = \hat{a}^\dagger \hat{a} + \hat{J}_z$; the field-ensemble and the inter-qubit coupling are represented by the constants λ and η, in that order.

This Dicke-LMG model, Eq.(1), may describe an experimentally feasible two-hyperfine-structure-defined-modes BEC coupled to a quantized cavity field mode through a one microwave photon process; *e.g.*, trapped hyperfine ground states of a Sodium BEC [12] inside a microwave cavity. It may also be possible to realize this model with arrays of interacting superconducting qubits [13] coupled to the quantized field mode of a coplanar waveguide resonator.

The ground state phase transition of the Dicke-LMG model has been studied in the thermodynamic limit, $N_a \to \infty$, within and without the rotating wave approximation (RWA)

[14] and in the quantum regime, using coherent states for both the field and ensemble, without the RWA [15]; these results show the existence of a finite size first order quantum phase transition and a second order super-radiant phase transition. Quantum phase transitions in the Dicke model have been associated with entanglement between the ensemble and the quantized field [16, 17] and bipartite entanglement among qubits due to finite-size effects [18–20].

In the following, we will show analytically, within the RWA, that the maximal entanglement, produced by the quantum phase transition, in the Lipkin-Meshkov-Glick model is retained, in the weak field-ensemble coupling regime, by the interacting qubit ensemble coupled to the quantized field. Furthermore, for a fixed field-ensemble coupling, λ, the inter-qubit coupling, η, defines a series of first order phase transitions related to the number of qubits in the ensemble, N_a, as the quotient $N_a/2$, while, for a fixed inter-qubit coupling, there exists one second order phase transition related to the couplings ratio [15] and a series of first order transitions similar to those in the Dicke model [18, 19]. Also, we will show analitically in the weak coupling regime, and numerically in general, that the inclusion of counter-rotating terms does not erase the possibility of obtaining a maximally entangled multi-qubit state.

The ground state of Eq.(1) has been approximated using the Holstein-Primakoff transformation and the boson expansion method. It was shown that a normal to superradiant phase transition occurs at zero temperature [14]; also, boson coherent states for the field and the Dicke basis for the qubit ensemble [15] have recently been used to show the existence of both a first order quantum phase transition and a second order superradiant phase transition. However, it is possible to find the exact ground state for the Hamiltonian in Eq.(1) in the RWA by partitioning the corresponding Hilbert space into subspaces with a mean total excitation number $\langle \hat{N} \rangle = n - N_a/2$ for $n = 0, 1, 2, \ldots$ due to conservation of the total number of excitations, $[\hat{H}, \hat{N}] = 0$. The ground state at the n-th subspace is given by $|\psi_g^{(n)}\rangle = \sum_{k=0}^{n} c_k^{(n)} |k\rangle_f |n-k-N_a/2\rangle$ such that $\hat{H}|\psi_g^{(n)}\rangle = E_g^{(n)}|\psi_g^{(n)}\rangle$ and $E_g^{(n)}$ is the lowest eigenvalue for the given subspace. The ket notation $|n\rangle_f$ refers to a Fock state of the field with n photons and the shorthand notation $|m\rangle \equiv |N_a/2, m\rangle$, $m = -N_a/2, 1 - N_a/2, \ldots, N_a/2 - 1, N_a/2$, is used for the Dicke state corresponding to the normalized superposition of all possible combinations of N_a atoms with $(N_a/2 + m)$ of them in the excited state and the rest, $(N_a/2 - m)$, in the ground state. The ground state is found as the lowest eigenvalue for the set $\{H^{(n)} = H_O^{(n)} + H_I^{(n)}\}$ of square matrices, with rank equal or less than $(N_a + 1)$, defined by the auxiliary matrices,

$$H_O^{(n)} = \omega_f \left(n - N_a/2\right) \mathbb{I}_{\tilde{n}}, \tag{2}$$

$$\left(H_I^{(n)}\right)_{i,j} = \delta_{i,j} d_i + \lambda N_a^{-1/2} \left(\delta_{i,j-1} o_j + \delta_{i,j+1} o_{j+1}\right), \tag{3}$$

where the identity matrix of rank d is given by $\mathbb{I}_d$, the row and column labels are in the range $i, j = \tilde{n}, \tilde{n} + 1, \ldots, n$, where $\tilde{n} = \max(0, n - N_a)$, for the photon number $n = 0, 1, 2, \ldots$ The symbol $\delta_{i,j}$ stands for Kronecker delta and the diagonal and off-diagonal terms are defined as,

$$d_j = (n - j - N_a/2) \left[\delta + \eta(n - j - N_a/2)/N_a\right], \tag{4}$$

$$o_j = \left[j(N_a + j - n)(n - j + 1)\right]^{1/2}. \tag{5}$$

The auxiliary matrix $H_I^{(n)}$ is a tri-diagonal symmetric real matrix with positive off-diagonal terms, *i.e.,* a Jacobi matrix, and its eigenvalues can be found analytically [21–24]. Furthermore, at the intersection of two ground state energies belonging to contiguous subspaces, a first order quantum phase transition is located.

The first ground state structure, which will be called vacuum phase from now on, corresponds to the vacuum field and the atomic ensemble ground state, $|\psi_g^{(0)}\rangle = |0\rangle| - N_a/2\rangle$. A first quantum phase transition, in a series of first order quantum phase transitions, is found at the critical coupling strength,

$$\lambda_{c1} = \left\{\left[\omega + (1/N_a - 1)\eta\right] \omega_f\right\}^{\frac{1}{2}}, \tag{6}$$

with the corresponding first non-vacuum state,

$$|\psi_g^{(1)}\rangle = c_0^{(1)} |0\rangle \, |1 - N_a/2\rangle + c_1^{(1)} |1\rangle \, |{-N_a/2}\rangle. \tag{7}$$

The amplitudes are given by the expressions $c_0^{(1)} = h/(h^2 + 1)^{1/2}$ and $c_1^{(1)} = 1/(h^2 + 1)^{1/2}$ related to the amplitude parameter,

$$h = \frac{(1 - 1/N_a)\eta + \delta - \{4\lambda^2 + [(1 - 1/N_a)\eta - \delta]^2\}^{1/2}}{2\lambda}. \tag{8}$$

Any extended Dicke model that conserves the total number of excitations has a first non-vacuum phase of the form given by Eq.(7), which includes the ground state of the atomic ensemble in the maximally entangled W-state $|1 - N_a/2\rangle$. The studied model, in the limit

$h \to 0$ yields amplitudes $c_0^{(1)} \to 0$ and $c_1^{(1)} \to 1$; *i.e.*, the state is the fully separable state $|1\rangle\,|-N_a/2\rangle$. In the opposite case, $h \to \infty$, the amplitudes are $c_0^{(1)} \to 1$ and $c_1^{(1)} \to 0$ and the state is the product of the vacuum field and the maximally entangled ensemble W-state, $|0\rangle\,|1 - N_a/2\rangle$. In the intermediate cases, $h \in (0, \infty)$, there is a finite probability of finding the ground state for this first non-vacuum phase in the maximally entangled W-state. Note that the case $h \to \infty$ can be obtained for small coupling strengths, $\lambda \to 0$; *i.e.*, the maximally entangled ground-state in the studied model is produced by the intra-ensemble interaction as expected by the fact that null field-ensemble coupling, $\lambda = 0$, delivers the LMG model.

In the absence of intra-ensemble interaction, $\eta = 0$, the Dicke model is obtained. It is well known that the shared bipartite concurrence at the first of a sequence of first order quantum phase transitions in the Dicke model is not maximal, nor is it at the subsequent first order quantum phase transitions [18]. Other extended Dicke models also lack the characteristic of a maximally entangled ground state at any first order quantum phase transition as well; *e.g.*, that includes a quantized field non-linearity [20].

In order to find out if the maximally entangled W-state shown at the first non-vacuum phase survives the inclusion of counter-rotating terms, it is possible to use the unitary transformation,

$$\hat{U} = e^{-\imath\xi(\hat{a}+\hat{a}^\dagger)\hat{J}_y}, \quad \xi = 2\lambda N_a^{-1/2}/(\omega + \omega_f). \tag{9}$$

In the weak coupling regime, $\lambda \ll \omega$ such that $\xi \ll 1$, and requiring a weak intra-ensemble interaction, $\eta \ll \omega$, it is possible to approximate,

$$\begin{aligned} \tilde{H}_{CR} &= \hat{U}^{-1}\hat{H}_{CR}\hat{U}, \\ &\approx \omega_f \hat{a}^\dagger\hat{a} + \omega\hat{J}_z + \eta N_a^{-1}\hat{J}_z^2 + \tilde{\lambda} N_a^{-1/2}(\hat{a}\hat{J}_+ + \hat{a}^\dagger\hat{J}_-), \end{aligned} \tag{10}$$

where the auxiliary coupling strength is given by $\tilde{\lambda} = 2\omega_f\lambda/(\omega + \omega_f)$. This is the original Hamiltonian with a modified coupling strength, *i.e.* the unitary transformation under weak coupling and intra-ensemble interaction acts as a rotating wave approximation, which has a first order quantum phase transition at the critical value on the weak coupling regime given by the expression,

$$\lambda_{c1}^{(CR)} = \frac{(\omega + \omega_f)}{2}\left[\frac{\omega + (1/N_a - 1)\eta}{\omega_f}\right]^{\frac{1}{2}}. \tag{11}$$

Note that on resonance the expression for the first critical coupling in the weak coupling regime, Eq.(11), is equal to the critical coupling in the rotating wave approximation, Eq.(6).

The ground state at the first non-vacuum phase is described again by Eq.(7) substituting $\lambda \to \tilde{\lambda}$. The maximally entangled W-state survives the inclusion of counter-rotating terms for the region of interest; *i.e.*, the weak coupling regime.

In the strong coupling regime, $\lambda \gg \omega$, it is possible to suppose the state of the system as the superposition of coherent states, $\left|\alpha\right\rangle_f \left|\theta,\phi\right\rangle_a$, with $\left|\theta,\phi\right\rangle_a = e^{-\imath \hat{J}_z \phi} e^{-\imath \hat{J}_y \theta} \left|N_a/2\right\rangle$. Under this representation [25],

$$
\begin{aligned}
\langle \hat{J}_z \rangle &= \frac{\sqrt{N_a}}{2} \sin\theta, && (12)\\
\langle \hat{J}_\pm \rangle &= \left[\frac{N_a}{2}\left(\frac{N_a}{2}+1\right) - \frac{N_a}{4}\sin^2\theta \pm \frac{\sqrt{N_a}}{2}\sin\theta \right]^{\frac{1}{2}} \\
&\quad \times \left[B_{N_a/2}(\theta) e^{\pm\imath\phi} + \left(\frac{\sin\theta}{2}\right)^{N_a} \right], && (13)
\end{aligned}
$$

with the coefficient

$$
\begin{aligned}
B_j(\beta) = \sum_m & \begin{pmatrix} 2j \\ j+m \end{pmatrix}^{\frac{1}{2}} \begin{pmatrix} 2j \\ j+m+1 \end{pmatrix}^{\frac{1}{2}} \\
& \times \left(\cos\frac{\beta}{2}\right)^{2j+2m+1} \left(\sin\frac{\beta}{2}\right)^{2j-2m-1} . \qquad (14)
\end{aligned}
$$

By minimizing the mean value energy for the three coherent parameters, $\{\alpha, \theta, \phi\}$, one can find the critical coupling in the strong limit; but this goes beyond the scope of this manuscript.

It is known that finite size and classical limit analyses, *i.e.* an infinitely large ensemble interacting with a coherent field, for Hamiltonians related to Dicke model yield identical critical couplings [18–20]. In order to round up the given analysis, we calculate the phase transition in the large number of particles limit. By using the Holstein-Primakoff transformation [26], $\hat{J}_z = \hat{b}^\dagger \hat{b} - N_a/2$, $\hat{J}_+ = N_a^{1/2} \hat{b}^\dagger \left(1 - \hat{b}^\dagger \hat{b}/N_a\right)^{1/2}$, $\hat{J}_+ = \hat{J}_-^\dagger$, and a Taylor expansion of the square root in terms of $(1/N_a)$, up to first order in $(1/N_a)$, $\left(1 - \hat{b}^\dagger \hat{b}/N_a\right)^{1/2} \approx 1 - \hat{b}^\dagger \hat{b}/2N_a$, up to a constant energy term of $N_a(\eta N_a - 2\omega)/2$, the Hamiltonian in Eq.(1), within the RWA, transforms into the expression,

$$
\begin{aligned}
\hat{H}_{CL} \approx\; & \omega_f \hat{a}^\dagger \hat{a} + (\omega - \eta N_a)\hat{b}^\dagger \hat{b} + \eta(\hat{b}^\dagger \hat{b})^2 \\
& + \lambda \left[\hat{a}\hat{b}^\dagger \left(1 - \frac{\hat{b}^\dagger \hat{b}}{2N_a}\right) + \left(1 - \frac{\hat{b}^\dagger \hat{b}}{2N_a}\right) \hat{a}^\dagger \hat{b} \right] . \qquad (15)
\end{aligned}
$$

In a Fock basis, the system presents a sequence of first order quantum phase transitions. In order to find the first of these, the problem transforms into diagonalizing the Hamiltonian in the one excitation subspace spanned by the states $\{|0,1\rangle, |1,0\rangle\}$. Thus, the ground state energy in the classical regime for the first non-vacuum phase is given by the expression,

$$E_{CLg}^{(1)} = \frac{1}{2}\Big[\omega + \omega_f + (1-N_a)\eta - \big\{4\lambda^2 + [\omega - \omega_f + (1/N_a - 1)\eta]^2\big\}^{\frac{1}{2}}\Big]. \tag{16}$$

The first critical coupling can be found for $E_g^{(1)} = 0$ and yields a form identical to the exact first critical coupling strength found above in Eq.(6),

$$\lambda_{c1}^{(CL)} = \{[\omega + (1/N_a - 1)\eta]\,\omega_f\}^{\frac{1}{2}}. \tag{17}$$

Equation (1), by using the Holstein-Primakoff transformation and the expansion of the square root the Hamiltonian, as stated above, transforms into the following Hamiltonian,

$$\begin{aligned}\hat{H}_{CLCR} \approx\ & \omega_f \hat{a}^\dagger \hat{a} + (\omega - \eta)\hat{b}^\dagger\hat{b} + \eta N_a^{-1}(\hat{b}^\dagger\hat{b})^2 + \lambda\left(\hat{a} + \hat{a}^\dagger\right)\left(\hat{b} + \hat{b}^\dagger\right) \\ & -\lambda(2N_a)^{-1/2}\left(\hat{a} + \hat{a}^\dagger\right)\hat{b}^\dagger\left(\hat{b} + \hat{b}^\dagger\right)\hat{b}.\end{aligned} \tag{18}$$

In order to find the phase transition, it is possible to use the coherent state basis $|\alpha\rangle_a |\beta\rangle_b$, *i.e.* a mean-field approximation, such that the mean value for the energy is given by

$$\begin{aligned}\langle \hat{H}_{CR}\rangle \approx\ & \omega_f |\alpha|^2 + (\omega - \eta)|\beta|^2 + \eta N^{-1}|\beta|^4 \\ & + \left(\lambda - \lambda(2N_a)^{-1}|\beta|^2\right)(\alpha + \alpha^*)(\beta + \beta^*)\end{aligned} \tag{19}$$

Optimizing with respect to the coherent parameters α and β, assuming them to be real numbers for the sake of simplicity, leads to a trivial solution and a non-trivial solution that requires $\lambda \geq [(\omega - \eta)\omega_f]^{1/2}/2$, thus a critical coupling strength for the Hamiltonian including counter rotating terms in the classical regime is found, as

$$\lambda_{c1}^{(CLCR)} = [(\omega - \eta)\,\omega_f]^{1/2}/2. \tag{20}$$

This critical coupling, $\lambda_{c1}^{(CLCR)}$, at which a second order superradiant phase transition occurs [15], is half of the critical strength found for the case without counter-rotating terms, $\lambda_{c1}^{(CL)}$, as $(1/N_a) \approx 0$ for $N \gg 2$. This is similar to what happens for the Dicke model in the classical limit result where accounting for counter-rotating terms halves the critical coupling found without the counter-rotating terms [27]. Also, this last result, $\lambda_{c1}^{(CLCR)}$, is directly related to that in the literature [14, 15].

In order to verify our analytical results, in Fig.(1), we show a numerical phase diagram defined by the coupling strength, λ, and interaction parameter, η, for a finite size qubit ensemble, $N_a = 5$. The maximum shared bipartite concurrence following the entangled web approach [4] and the field-ensemble entanglement probed through von Neumann entropy, $\langle \hat{S} \rangle = -\mathrm{Tr}_a \left(\rho_a \log \rho_a \right)$, also known as entropy of entanglement [2], of the reduced qubit ensemble obtained after tracing out the field from the total density matrix, $\rho_a = \mathrm{tr}_f \rho$, are shown in Fig.(1) for the case within the RWA and in Fig.(2) for the full model. The first and second critical couplings shown in dashed black lines in Fig.(1) are analytically exact obtained from the analysis presented above. As analytical and numerically exact results agree for this case, only the numerical graphics are shown. Likewise, in Fig.(2), by following an equivalent approach to that presented in reference [20], numerical results and the first critical coupling including counter-rotating terms are shown. Note that, for this case, it is not possible to obtain closed form exact results but the analytical critical coupling found for the weak coupling regime is in agreement with the results obtained numerically.

A maximally entangled ground state is found for very small values of the coupling strength, such that $\eta/\lambda \gg 1$. Stability analysis of the Hamiltonian in the classical limit show that a pitchfork bifurcation point of the classical dynamics may occur at these coupling strength ratios [28], as conjectured for coupled giant spins [29].

Figures 1(a) and 1(b) show that it is possible to have a W-ground-state for a pentapartite ensemble as expected in the weak coupling limit. For five qubits, there exists a large parameter region at the first non-vacuum phase where maximal value for the maximum shared bipartite concurrence, $C_W = 2/5$, is reached. This value wanes subsequently as expected. The entropy of entanglement rises to a maximum after the first zone, which is related to a second order phase transition, a behavior shared by all studied cases, $N_a \in [2, 6]$, to decay for very large values of the coupling strength, $\lambda \gg \omega$.

In contrast to the Dicke model, where the first critical coupling is independent of the ensemble size, in this extended Dicke model the ensemble size plays a role in the first critical coupling, *e.g.*, Eq.(6). As the involved term is of the form $-N^{-1}$, it stops playing an important role for large ensemble sizes. In the Dicke model, the vacuum phase remains fixed as the first critical coupling stays constant for an increasing number of atoms, while the phases beyond the vacuum phase reduce to allow just the vacuum and classical superradiant phases for an infinitely large atomic ensemble. In this extended Dicke model, the area of all the different phases reduces as the ensemble size increases.

In summary, we have analytically derived the exact critical coupling value and ground states at the first quantum phase transition for a BEC-cavity-QED system in the rotating wave approximation, including counter-rotating terms, and in the classical limit both excluding and including counter-rotating terms. Subsequent ground states of first order quantum phase transitions have been shown through analytical derivations and numerical simulations. The results obtained correlate among themselves in a similar way as the Dicke model.

While the exact critical coupling at the first of the quantum phase transitions connects with known mean-field results, our treatment allows to follow the quantum correlations between the field and ensemble as well as among ensemble components. Thus, a striking feature is revealed: the existence of a maximally entangled atomic ground state within the limits of a certain phase space region at the first non-vacuum phase. The parameter region where this maximally entangled W-state is found decreases as the ensemble grows in size. The existence of this maximal entangled qubit ensemble state survives even the addition of counter rotating terms. Numerical results for the maximum shared bipartite concurrence, following the entangled web approach, and the field-ensemble entanglement, probed through von Neuman entropy of the reduced two-level ensemble, were presented. Agreement between the exact analytical and the numerical results was obtained.

We believe that our results show that highly tunable experimental systems realizing an interacting ensemble of qubits driven by a quantized field in the weak coupling regime may be used as a source of entanglement for quantum information processing;*e.g.*, an array of interacting superconducting flux qubits coupled to a coplanar waveguide resonator following the building blocks presented in references [13, 30] or a two ground-state-hyperfine-structure-defined mode BEC driven by a single mode of microwave cavity building on known technology like that in Ref. [12], with the inconvenience that the center of mass motion may couple to the internal degrees of freedom [31].



Figure 1. (Color online) The phase diagram of the finite size extended Dicke Hamiltonian for the case of a penta-partite qubit ensemble, $N_a = 5$, in the parameter space defined by the field-ensemble coupling strength, λ, and the intra-ensemble interaction strength, η. (a) Maximum shared bipartite concurrence in the sense of entangled webs, C_w. (b) Entropy of entanglement, $\langle\hat{S}\rangle$, between the field and the qubit ensemble. The corresponding minima and maxima values for the color legend are shown below it. The dashed lines show the two first exact critical couplings.

---

[1] E. Schrödinger, Naturwissenschaften **23**, 807 (1935).

E. Schrödinger, Naturwissenschaften **23**, 823 (1935).

E. Schrödinger, Naturwissenschaften **23**, 844 (1935).

A. Einstein, B. Podolsky, and N. Rosen, Phys. Rev. **47**, 777 (1935).

J. S. Bell, Physics **1**, 195 (1964).

[2] M. Nielsen and I. Chuang, *Quantum Information and Computation* (Cambridge University Press, 2000).

J. Preskill, J. Mod. Opt. **47**, 127 (2000).

[3] W. Dür, G. Vidal, and J. I. Cirac, Phys. Rev. A **62**, 062314 (2000).

[4] M. Koashi, V. Bužek, and N. Imoto, Phys. Rev. A **62**, 050302(R) (2000).

[5] W. Dür, Phys. Rev. A **63**, 020303 (2001).

[6] H. J. Briegel and R. Raussendorf, Phys. Rev. Lett. **86**, 910 (2001).

[7] A. Sen(De), U. Sen, M. Wiesniak, D. Kaszlikowski, and M. Zukowski, Phys. Rev. A **68**, 062306



Figure 2. (Color online) The phase diagram of the finite size extended Dicke Hamiltonian including counter rotating terms for the case of a penta-partite qubit ensemble, $N_a = 5$, in the parameter space defined by the field-ensemble coupling strength, λ, and the intra-ensemble interaction strength, η. (a) Maximum shared bipartite concurrence in the sense of entangled webs, C_w. (b) Entropy of entanglement, $\langle\hat{S}\rangle$, between the field and the qubit ensemble. The corresponding minima and maxima values for the color legend are shown below it. The dashed line show the first exact critical coupling in the weak coupling regime.

(2003).

[8] H. J. Lipkin, N. Meshkov, and A. J. Glick, Nucl. Phys **62**, 188 (1965).
J. Vidal, G. Palacios, and R. Mosseri, Phys. Rev. A **69**, 022107 (2004).
J. Vidal, R. Mosseri, and J. Dukelsky, Phys. Rev. A **69**, 054101 (2004).

[9] J. Vidal, G. Palacios, and C. Aslangul, Phys. Rev. A **70**, 062304 (2004).

[10] S. Morrison and A. S. Parkins, Phys. Rev. Lett. **100**, 040403 (2008).

[11] J. Larson, Europhys. Lett. **90**, 54001 (2010).

[12] A. Görlitz, T. L. Gustavson, A. E. Leanhardt, R. Low, A. P. Chikkatur, S. Gupta, S. Inouye, D. E. Pritchard, and W. Ketterle, Phys. Rev. Lett. **90**, 090401 (2003).

[13] D. I. Tsomokos, S. Ashhab, and F. Nori, Phys. Rev. A **82**, 052311 (2010).

[14] G. Chen, Z. Chen, and J.-Q. Liang, Europhys. Lett. **80**, 40004 (2007).

[15] Q.-H. Chen, T. Liu, Y.-Y. Zhang, and K.-L. Wang, Phys. Rev. A **82**, 053841 (2010).

[16] N. Lambert, C. Emary, and T. Brandes, Phys. Rev. Lett. **92**, 073602 (2004).

[17] J. Vidal and S. Dusuel, Europhys. Lett. **74**, 817 (2006).

[18] V. Bužek, M. Orszag, and M. Roško, Phys. Rev. Lett. **94**, 163601 (2005).

[19] O. Tsyplyatyev and D. Loss, J. Phys.: Conf. Ser. **193**, 012134 (2009).

[20] B. M. Rodríguez-Lara and R.-K. Lee, J. Opt. Soc. Am. B **27**, 2443 (2010).

[21] J. D. Swalen and L. Pierce, J. Math. Phys. **2**, 736 (1961).

[22] L. Pierce, J. Math. Phys. **2**, 740 (1961).

[23] R. Haydock, V. Heine, and M. J. Kelly, J. Phys. C: Solid State Phys. **5**, 2845 (1972).

[24] H. A. Yamani and M. S. Abdelmonem, J. Phys. A: Math. Gen. **30**, 2889 (1997).

[25] G. Nienhuis and S. J. van Enk, Phys. Scr. **T48**, 87 (1993).

[26] T. Holstein and H. Primakoff, Phys. Rev. **58**, 1098 (1940).

[27] H. J. Carmichael, C. W. Gardiner, and D. F. Walls, Phys. Lett. **46**, 47 (1973).

[28] B. M. Rodríguez-Lara and R.-K. Lee, arxiv:1102.3747v1[quant-ph](2011).

[29] A. P. Hines, R. H. McKenzie, and G. J. Milburn, Phys. Rev. A **71**, 042303 (2005).

[30] J. Q. You and F. Nori, Phys. Rev. B **68**, 064509 (2003).

[31] J. Larson and M. Lewenstein, New. J. Phys **11**, 063027 (2009).

L. O. Castaños and R. Jáuregui, Phys. Rev. A **82**, 053815 (2010).